

WAVE EQUATION WITH THREE-INVERSE SQUARE POTENTIAL ON R_+^3

Yehdih Mohamed Abdelhay, Badahi Mohamed and Mohamed Vall Ould Moustapha

Abstract In this note we give explicit solutions to the wave equation associated to the Schrödinger operator with three-inverse square potential on R_+^3 .

Key words : Three inverse square potential, Cauchy problem, Wave equation, Lauricella hypergeometric functions.

1 Introduction and statement of results

The wave equation, the heat equation and the Laplace equation are known as three fundamental equations in partial differential equations and occur in many branches of physics, in applied mathematics and in engineering. In this note we give explicit formulas for the solutions of the following Cauchy problem for the wave equation with three-inverse square potential

$$(W)_{(\nu, \nu', \nu'')} \begin{cases} (a) & [\Delta + v_{\nu, \nu', \nu''}(x)]u(t, p) = \frac{\partial^2}{\partial t^2} u(t, p) \\ (b) & u(0, p) = 0 \end{cases} \quad \begin{matrix} (t, p) \in R \times R_+^3 \\ \frac{\partial}{\partial t} u(0, p) = u_1(p), u_1 \in C_0^\infty(R_+^3) \end{matrix}$$

with $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ is the Laplacien of R^3 , the three inverse square potential is given by $v_{\nu, \nu', \nu''}(x, y, z) = \frac{1/4-\nu^2}{x^2} + \frac{1/4-\nu'^2}{y^2} + \frac{1/4-\nu''^2}{z^2}$ where ν, ν', ν'' are real parameters. The Cauchy problem for the wave equation with the inverse square potential in Euclidean space \mathbb{R}^n is extensively studied (Cheeger and Taylor [3]). The bi-inverse square potential has been considered by (Boyer [2]) and Ould Moustapha [7]. The case considered most frequently is obviously the one where $(\nu, \nu', \nu'') = (\pm 1/2, \pm 1/2, \pm 1/2)$, the equation in $(W)_{\nu, \nu', \nu''}$ then turns into the classical wave equation on the Euclidean space \mathbb{R}^3 and this equation appears in several branches of mathematics and physics (Folland [5], p.171). Now we state the main results of this paper:

Theorem A For $(t, p, p') \in R_+ \times \mathbb{R}_+^3 \times \mathbb{R}_+^3$ the functions:

$$W_{(b, b', b'')}(t, p, p') = \frac{c_3 (xx')^b (yy')^{b'} (zz')^{b''}}{(t^2 - |p+p'|^2)^{1+b+b'+b''}} F_A^{(3)} \left(1+b+b'+b'', b, b', 2b, 2b', 2b'', \frac{-4xx'}{t^2 - |p+p'|^2}, \frac{-4yy'}{t^2 - |p+p'|^2}, \frac{-4zz'}{t^2 - |p+p'|^2} \right)$$

with $b \in \{\beta, 1-\beta\}$, $b' \in \{\beta', 1-\beta'\}$, $b'' \in \{\beta'', 1-\beta''\}$ are independent solutions of the wave equation with three-inverse square potential on \mathbb{R}_+^3 (a) where $\beta = 1/2 + \nu$ and $\beta' = 1/2 + \nu'$, $\beta'' = 1/2 + \nu''$ and $F_A^{(3)}(a, b, b', b'', c, c', c''; w, w', w'')$ is the three variables triple series $F_A^{(3)}$ Lauricella hypergeometric function given by [1], p.114

$$F_A^{(3)}(a, b, b', b'', c, c', c''; w, w', w'') = \sum_{m, n, p \geq 0} \frac{(a)_{m+n+p} (b)_m (b')_n (b'')_p}{(c)_m m! (c')_n n! (c'')_p p!} w^m w'^n w''^p \quad (1.1)$$

Theorem B The Cauchy problem for the wave equation with three-inverse square potential on the \mathbb{R}_+^3 has the solutions given by:

$$u(t, p) = \frac{\partial}{\partial t} \int_{|p+p'| < t} W_{(b, b', b'')}(t, p, p') f(p') dp' \quad (1.2)$$

where the kernel $W_{(b, b', b'')}$ is as in the Theorem A and the constant c_3 is given by

$$c_3 = \frac{(-1)^{b+b'+b''} \Gamma(1/2 + b + b' + b'')}{\sqrt{\pi} \Gamma(1/2 + b) \Gamma(1/2 + b') \Gamma(1/2 + b'')} \quad (1.3)$$

with $dp' = dx' dy' dz'$ is the Lebesgue measure on \mathbb{R}^3

2 Wave equation with three-inverse square potential on Euclidean space \mathbb{R}^3

Proof of Theorem A

In what follows we give a direct proof of the theorem A.

Let $a = t^2 - |p + p'|^2$, $t \in \mathbb{R}$, $p, p' \in \mathbb{R}^{*3}$ set:

$$\Omega\varphi(t, p) = (xx')^{-\beta} (yy')^{-\beta'} (zz')^{-\beta''} a^{-\alpha} \times$$

$$\left[\Delta - \frac{\beta(\beta-1)}{x^2} - \frac{\beta'(\beta'-1)}{y^2} - \frac{\beta''(\beta''-1)}{z^2} - \frac{\partial^2}{\partial t^2} \right] (xx')^\beta (yy')^{\beta'} (zz')^{\beta''} a^\alpha \varphi(t, x) \quad (2.1)$$

then we have

$$\begin{aligned} \Omega\varphi(t, p) = & \left\{ \Delta - \frac{\partial^2}{\partial t^2} + \left[\frac{2\beta}{x} - \frac{4\alpha(x+x')}{a} \right] \frac{\partial}{\partial x} + \left[\frac{2\beta'}{y} - \frac{4\alpha(y+y')}{a} \right] \frac{\partial}{\partial y} + \left[\frac{2\beta''}{z} - \frac{4\alpha(z+z')}{a} \right] \frac{\partial}{\partial z} - \frac{4\alpha t}{a} \frac{\partial}{\partial t} \right. \\ & \left. - \frac{4\alpha}{a} \left[\frac{\beta x'}{x} + \frac{\beta' y'}{y} + \frac{\beta'' z'}{z} \right] - \frac{4\alpha}{a} [\alpha + 1 + \beta + \beta' + \beta''] \right\} \varphi(t, x) \end{aligned} \quad (2.2)$$

Now set

$$w = \frac{-4xx'}{t^2 - |p + p'|^2}, w' = \frac{-4yy'}{t^2 - |p + p'|^2}, w'' = \frac{-4zz'}{t^2 - |p + p'|^2} \quad (2.3)$$

we can write:

$$\frac{\partial}{\partial x} = \frac{\partial w}{\partial x} \frac{\partial}{\partial w} + \frac{\partial w'}{\partial x} \frac{\partial}{\partial w'} + \frac{\partial w''}{\partial x} \frac{\partial}{\partial w''}; \quad \frac{\partial}{\partial y} = \frac{\partial w}{\partial y} \frac{\partial}{\partial w} + \frac{\partial w'}{\partial y} \frac{\partial}{\partial w'} + \frac{\partial w''}{\partial y} \frac{\partial}{\partial w''} \quad (2.4)$$

$$\frac{\partial}{\partial z} = \frac{\partial w}{\partial z} \frac{\partial}{\partial w} + \frac{\partial w'}{\partial z} \frac{\partial}{\partial w'} + \frac{\partial w''}{\partial z} \frac{\partial}{\partial w''}; \quad \frac{\partial}{\partial t} = \frac{\partial w}{\partial t} \frac{\partial}{\partial w} + \frac{\partial w'}{\partial t} \frac{\partial}{\partial w'} + \frac{\partial w''}{\partial t} \frac{\partial}{\partial w''} \quad (2.5)$$

We have:

$$\begin{aligned} \Omega\varphi(t, p) = & \left[\left(\frac{\partial w}{\partial x} \right)^2 + \left(\frac{\partial w}{\partial y} \right)^2 + \left(\frac{\partial w}{\partial z} \right)^2 - \left(\frac{\partial w}{\partial t} \right)^2 \right] \frac{\partial^2}{\partial w^2} \\ & + \left[\left(\frac{\partial w'}{\partial x} \right)^2 + \left(\frac{\partial w'}{\partial y} \right)^2 + \left(\frac{\partial w'}{\partial z} \right)^2 - \left(\frac{\partial w'}{\partial t} \right)^2 \right] \frac{\partial^2}{\partial w'^2} + \left[\left(\frac{\partial w''}{\partial x} \right)^2 + \left(\frac{\partial w''}{\partial y} \right)^2 + \left(\frac{\partial w''}{\partial z} \right)^2 - \left(\frac{\partial w''}{\partial t} \right)^2 \right] \frac{\partial^2}{\partial w''^2} \\ & + 2 \left[\frac{\partial w}{\partial x} \frac{\partial w'}{\partial x} + \frac{\partial w}{\partial y} \frac{\partial w'}{\partial y} + \frac{\partial w}{\partial z} \frac{\partial w'}{\partial z} - \frac{\partial w}{\partial t} \frac{\partial w'}{\partial t} \right] \frac{\partial^2}{\partial w \partial w'} + 2 \left[\frac{\partial w}{\partial x} \frac{\partial w''}{\partial x} + \frac{\partial w}{\partial y} \frac{\partial w''}{\partial y} + \frac{\partial w}{\partial z} \frac{\partial w''}{\partial z} - \frac{\partial w}{\partial t} \frac{\partial w''}{\partial t} \right] \frac{\partial^2}{\partial w \partial w''} \\ & + 2 \left[\frac{\partial w'}{\partial x} \frac{\partial w''}{\partial x} + \frac{\partial w'}{\partial y} \frac{\partial w''}{\partial y} + \frac{\partial w'}{\partial z} \frac{\partial w''}{\partial z} - \frac{\partial w'}{\partial t} \frac{\partial w''}{\partial t} \right] \frac{\partial^2}{\partial w' \partial w''} + \left[\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} - \frac{\partial^2 w}{\partial t^2} \right] \frac{\partial}{\partial w} \\ & + \left[\frac{\partial^2 w'}{\partial x^2} + \frac{\partial^2 w'}{\partial y^2} + \frac{\partial^2 w'}{\partial z^2} - \frac{\partial^2 w'}{\partial t^2} \right] \frac{\partial}{\partial w'} + \left[\frac{\partial^2 w''}{\partial x^2} + \frac{\partial^2 w''}{\partial y^2} + \frac{\partial^2 w''}{\partial z^2} - \frac{\partial^2 w''}{\partial t^2} \right] \frac{\partial}{\partial w''} \\ & + [A_x \frac{\partial w}{\partial x} + A_y \frac{\partial w}{\partial y} + A_z \frac{\partial w}{\partial z} - A_t \frac{\partial w}{\partial t}] \frac{\partial}{\partial w} + [A_x \frac{\partial w'}{\partial x} + A_y \frac{\partial w'}{\partial y} + A_z \frac{\partial w'}{\partial z} - A_t \frac{\partial w'}{\partial t}] \frac{\partial}{\partial w'} \\ & + [A_x \frac{\partial w''}{\partial x} + A_y \frac{\partial w''}{\partial y} + A_z \frac{\partial w''}{\partial z} - A_t \frac{\partial w''}{\partial t}] \frac{\partial}{\partial w''} + \frac{4\alpha}{a} \left[\frac{\beta x'}{x} + \frac{\beta' y'}{y} + \frac{\beta'' z'}{z} \right] + \frac{4\alpha}{a} [\alpha + 1 + \beta + \beta' + \beta''] \varphi(z, z') \end{aligned} \quad (2.6)$$

where

$$A_x = \frac{2\beta}{x} - \frac{4\alpha(x+x')}{a}, A_y = \frac{2\beta'}{y} - \frac{4\alpha(y+y')}{a}, A_z = \frac{2\beta''}{z} - \frac{4\alpha(z+z')}{a}, A_t = \frac{4\alpha t}{a} \quad (2.7)$$

We have:

$$\frac{\partial w}{\partial x} = \frac{-4x'a - 8(x+x')xx'}{a^2}; \quad \frac{\partial w'}{\partial x} = \frac{-8yy'(x+x')}{a^2}; \quad \frac{\partial w''}{\partial x} = \frac{-8zz'(x+x')}{a^2} \quad (2.8)$$

$$\frac{\partial w}{\partial y} = \frac{-8xx'(y+y')}{a^2}; \quad \frac{\partial w'}{\partial y} = \frac{-4y'a - 8yy'(y+y')}{a^2}; \quad \frac{\partial w''}{\partial y} = \frac{-8zz'(y+y')}{a^2} \quad (2.9)$$

$$\frac{\partial w}{\partial z} = \frac{-8xx'(z+z')}{a^2}; \quad \frac{\partial w'}{\partial z} = \frac{-8yy'(z+z')}{a^2}; \quad \frac{\partial w''}{\partial z} = \frac{-4z'a - 8zz'(z+z')}{a^2} \quad (2.10)$$

$$\frac{\partial w}{\partial t} = \frac{8xx't}{a^2}; \quad \frac{\partial w'}{\partial t} = \frac{8yy't}{a^2}; \quad \frac{\partial w''}{\partial t} = \frac{8zz't}{a^2} \quad (2.11)$$

$$\frac{\partial^2 w}{\partial x^2} = \frac{-24xx'a^2 - 16x'^2a^2 - 32(x+x')^2xx'a}{a^4}; \quad \frac{\partial^2 w}{\partial y^2} = \frac{-8yy'a^2 - 32xx'(y+y')^2a}{a^4}; \quad \frac{\partial^2 w}{\partial z^2} = \frac{-8zz'a^2 - 32xx'(z+z')^2a}{a^4} \quad (2.12)$$

$$\frac{\partial^2 w'}{\partial x^2} = \frac{-8xx'a^2 - 32yy'(x+x')^2a}{a^4}; \quad \frac{\partial^2 w'}{\partial y^2} = \frac{-24yy'a^2 - 16y'^2a^2 - 32(y+y')^2yy'a}{a^4}; \quad \frac{\partial^2 w'}{\partial z^2} = \frac{-8yy'a^2 - 32azz'(y+y')^2a}{a^4} \quad (2.13)$$

$$\frac{\partial^2 w''}{\partial x^2} = \frac{-8xx'a^2 - 32zz'(x+x')^2a}{a^4}; \quad \frac{\partial^2 w''}{\partial y^2} = \frac{-8yy'a^2 - 32zz'(y+y')^2a}{a^4}; \quad \frac{\partial^2 w''}{\partial z^2} = \frac{-24zz'a^2 - 16z'^2a^2 - 32(z+z')^2zz'a}{a^4} \quad (2.14)$$

$$\frac{\partial^2 w}{\partial t^2} = \frac{8xx'a^2 - 32axx't^2}{a^4}; \quad \frac{\partial^2 w'}{\partial t^2} = \frac{8yy'a^2 - 32ayy't^2}{a^4}; \quad \frac{\partial^2 w''}{\partial t^2} = \frac{8zz'a^2 - 32azz't^2}{a^4} \quad (2.15)$$

from (2.8) – (2.11) we have

$$\left(\frac{\partial w}{\partial x} \right)^2 + \left(\frac{\partial w}{\partial y} \right)^2 + \left(\frac{\partial w}{\partial z} \right)^2 - \left(\frac{\partial w}{\partial t} \right)^2 = \frac{w^2}{x^2} (1 - w) \quad (2.16)$$

$$\left(\frac{\partial w'}{\partial x} \right)^2 + \left(\frac{\partial w'}{\partial y} \right)^2 + \left(\frac{\partial w'}{\partial z} \right)^2 - \left(\frac{\partial w'}{\partial t} \right)^2 = \frac{w'^2}{y^2} (1 - w') \quad (2.17)$$

$$\left(\frac{\partial w''}{\partial x}\right)^2 + \left(\frac{\partial w''}{\partial y}\right)^2 + \left(\frac{\partial w''}{\partial z}\right)^2 - \left(\frac{\partial w''}{\partial t}\right)^2 = \frac{w''^2}{z^2}(1 - w'') \quad (2.18)$$

$$2 \left[\frac{\partial w}{\partial x} \frac{\partial w'}{\partial x} + \frac{\partial w}{\partial y} \frac{\partial w'}{\partial y} + \frac{\partial w}{\partial z} \frac{\partial w'}{\partial z} - \frac{\partial w}{\partial t} \frac{\partial w'}{\partial t} \right] = \frac{w^2}{x^2} w' + \frac{w'^2}{y^2} w \quad (2.19)$$

$$2 \left[\frac{\partial w}{\partial x} \frac{\partial w''}{\partial x} + \frac{\partial w}{\partial y} \frac{\partial w''}{\partial y} + \frac{\partial w}{\partial z} \frac{\partial w''}{\partial z} - \frac{\partial w}{\partial t} \frac{\partial w''}{\partial t} \right] = \frac{w^2}{x^2} w'' + \frac{w''^2}{z^2} w \quad (2.20)$$

$$2 \left[\frac{\partial w'}{\partial x} \frac{\partial w''}{\partial x} + \frac{\partial w'}{\partial y} \frac{\partial w''}{\partial y} + \frac{\partial w'}{\partial z} \frac{\partial w''}{\partial z} - \frac{\partial w'}{\partial t} \frac{\partial w''}{\partial t} \right] = \frac{w'^2}{y^2} w'' + \frac{w''^2}{z^2} w' \quad (2.21)$$

from (2.12) – (2.15) we have

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} - \frac{\partial^2 w}{\partial t^2} = \frac{-w^2}{x^2} + \frac{2}{a} w \quad (2.22)$$

$$\frac{\partial^2 w'}{\partial x^2} + \frac{\partial^2 w'}{\partial y^2} + \frac{\partial^2 w'}{\partial z^2} - \frac{\partial^2 w'}{\partial t^2} = \frac{-w'^2}{y^2} + \frac{2}{a} w' \quad (2.23)$$

$$\frac{\partial^2 w''}{\partial x^2} + \frac{\partial^2 w''}{\partial y^2} + \frac{\partial^2 w''}{\partial z^2} - \frac{\partial^2 w''}{\partial t^2} = \frac{-w''^2}{z^2} + \frac{2}{a} w'' \quad (2.24)$$

from (2.7) – (2.11) we have

$$A_x \frac{\partial w}{\partial x} + A_y \frac{\partial w}{\partial y} + A_z \frac{\partial w}{\partial z} - A_t \frac{\partial w}{\partial t} = \frac{4\alpha + 4(\beta + \beta' + \beta'')}{a} w + 2\beta \frac{w}{x^2} + \alpha \frac{w^2}{x^2} - \beta \frac{w^2}{x^2} + \left(\frac{\beta' w'}{y^2} + \frac{\beta'' w''}{z^2} \right) w \quad (2.25)$$

$$A_x \frac{\partial w'}{\partial x} + A_y \frac{\partial w'}{\partial y} + A_z \frac{\partial w'}{\partial z} - A_t \frac{\partial w'}{\partial t} = \frac{4\alpha + 4(\beta + \beta' + \beta'')}{a} w' + 2\beta \frac{w'}{y^2} + \alpha \frac{w'^2}{y^2} - \beta \frac{w'^2}{y^2} + \left(\frac{\beta w}{x^2} + \frac{\beta'' w''}{z^2} \right) w' \quad (2.26)$$

$$A_x \frac{\partial w''}{\partial x} + A_y \frac{\partial w''}{\partial y} + A_z \frac{\partial w''}{\partial z} - A_t \frac{\partial w''}{\partial t} = \frac{4\alpha + 4(\beta + \beta' + \beta'')}{a} w'' + 2\beta'' \frac{w''}{z^2} + \alpha \frac{w''^2}{z^2} - \beta'' \frac{w''^2}{z^2} + \left(\frac{\beta w}{x^2} + \frac{\beta' w'}{y^2} \right) w'' \quad (2.27)$$

To replace in the formula (2.6) using the formulas (2.16) – (2.27) we get:

$$\Omega\varphi = wx^{-2} A_{\alpha,\beta}(w, w', w'')\varphi + w'y^{-2} A_{\alpha,\beta'}(w', w, w'')\varphi + w''z^{-2} A_{\alpha,\beta''}(w'', w, w')\varphi +$$

$$\frac{4}{a}(\alpha + \beta + \beta' + \beta'' + 1) \left[\left(w \frac{\partial}{\partial w} + w' \frac{\partial}{\partial w'} + w'' \frac{\partial}{\partial w''} \right) \varphi(w, w', w'') - \varphi(w, w', w'') \right] \quad (2.28)$$

Take $\alpha = -1 - \beta - \beta' - \beta''$ we get $\Omega\varphi = 0$ is equivalent to

$$wx^{-2} A_{\alpha,\beta}(w, w', w'')\varphi + w'y^{-2} A_{\alpha,\beta'}(w', w, w'')\varphi + w''z^{-2} A_{\alpha,\beta''}(w'', w, w')\varphi = 0 \quad (2.29)$$

with

$$A_{\alpha,\beta}(w, w', w'')\varphi(w, w', w'') = [w(1-w) \frac{\partial^2}{\partial w^2} - w(w' \frac{\partial^2}{\partial w' \partial w} + w'' \frac{\partial^2}{\partial w'' \partial w}) + \\ + [2\beta + (-\alpha + \beta + 1)w] \frac{\partial}{\partial w} - \beta' w' \frac{\partial}{\partial w'} - \beta'' w'' \frac{\partial}{\partial w''} + \alpha\beta] \varphi(w, w', w'') \quad (2.30)$$

From the formula (2.29) we have

$$A_{\alpha,\beta}(w, w', w'')\varphi(w, w', w'') = A_{\alpha,\beta'}(w', w, w'')\varphi(w, w', w'') = A_{\alpha,\beta''}(w'', w, w')\varphi(w, w', w'') = 0 \quad (2.31)$$

$$[w(1-w) \frac{\partial^2}{\partial w^2} - w(w' \frac{\partial^2}{\partial w' \partial w} + w'' \frac{\partial^2}{\partial w'' \partial w}) + [2\beta + (-\alpha + \beta + 1)w] \frac{\partial}{\partial w} - \beta' w' \frac{\partial}{\partial w'} - \beta'' w'' \frac{\partial}{\partial w''} + \alpha\beta] \varphi(w, w', w'') = 0,$$

$$[w'(1-w') \frac{\partial^2}{\partial w'^2} - w'(w \frac{\partial^2}{\partial w \partial w'} + w'' \frac{\partial^2}{\partial w'' \partial w'}) + [2\beta' + (-\alpha + \beta' + 1)w'] \frac{\partial}{\partial w'} - \beta w \frac{\partial}{\partial w} - \beta'' w'' \frac{\partial}{\partial w''} + \alpha\beta'] \varphi(w, w', w'') = 0,$$

$$[w''(1-w'') \frac{\partial^2}{\partial w''^2} - w''(w' \frac{\partial^2}{\partial w' \partial w''} + w \frac{\partial^2}{\partial w \partial w''}) + [2\beta'' + (-\alpha + \beta'' + 1)w''] \frac{\partial}{\partial w''} - \beta w \frac{\partial}{\partial w} - \beta' w' \frac{\partial}{\partial w'} + \alpha\beta''] \varphi(w, w', w'') = 0,$$

This is an $F_A^{(3)}$ three variable Lauricella hypergeometric system and for $2\beta \neq 1$ and $2\beta' \neq 1$ $2\beta'' \neq 1$ the system has six independent solutions of the form [4], p.150 – 151:

- $F_A^{(3)}(-\alpha, \beta, \beta', \beta'', 2\beta, 2\beta', 2\beta'', w, w', w'')$,
- $w^{1-2\beta} F_A^{(3)}(-\alpha + 1 - 2\beta, 1 - \beta, \beta', \beta'', 2 - 2\beta, 2\beta', 2\beta'', w, w', w'')$,

- $w'^{1-2\beta'} F_A^{(3)}(-\alpha+1-2\beta', \beta, 1-\beta', \beta'', 2\beta, 1-2\beta', 2\beta'', w, w', w''),$
- $w''^{1-2\beta''} F_A^{(3)}(-\alpha+1-2\beta'', \beta, \beta', 1-\beta'', 2\beta, 2\beta', 2-2\beta'', w, w', w''),$
- $w^{1-2\beta} w'^{1-2\beta'} F_A^{(3)}(-\alpha+2-2\beta-2\beta', 1-\beta, 1-\beta', \beta'', 2-2\beta, 2-2\beta', 2\beta'', w, w', w''),$
- $w^{1-2\beta} w''^{1-2\beta''} F_A^{(3)}(-\alpha+2-2\beta-2\beta'', 1-\beta, \beta', 1-\beta'', 2-2\beta, 2\beta', 2-2\beta'', w, w', w''),$
- $w'^{1-2\beta} w''^{1-2\beta} F_A^{(3)}(-\alpha+2-2\beta'-2\beta'', \beta, 1-\beta', 1-\beta'', 2\beta, 2-2\beta', 2-2\beta'', w, w', w''),$
- $w^{1-2\beta} w'^{1-2\beta} w''^{1-2\beta} F_A^{(3)}(-\alpha+3-2\beta-2\beta'-2\beta'', 1-\beta, 1-\beta', 1-\beta'', 2-2\beta, 2-2\beta', 2-2\beta'', w, w', w'')$

And the proof of the theorem 1.1 is finished.

3 Cauchy problem for the wave equation with the three-inverse square potential on R_+^3

Proof of the Theorem B

Lemma 3.1 Let $F_A^{(3)}$ be the Appell hypergeometric function with $(h, k, l) \in R^3$ and $a \in R^*$ then we have:

i)

$$\frac{d}{da} \left[a^\alpha F_A^{(3)}(-\alpha, \beta, \beta', \beta'', 2\beta, 2\beta', 2\beta'', h/a, k/a, l/a) \right] = -\alpha a^{\alpha-1} \times$$

$$F_A^{(3)}(-\alpha+1, \beta, \beta', \beta'', 2\beta, 2\beta', 2\beta'', h/a, k/a, l/a) \quad (3.1)$$

ii)

$$a^\alpha \Gamma(-\alpha) F_A^{(3)}(-\alpha, \beta, \beta', \beta'', 2\beta, 2\beta', 2\beta'', h/a, k/a, l/a) = \Gamma(\beta+1/2) \Gamma(\beta'+1/2) \Gamma(\beta''+1/2) \times$$

$$\int_0^\infty e^{-(a-h/2-k/2-l/2)t} t^{-\alpha-1} \left(\frac{th}{4} \right)^{1/2-\beta} \left(\frac{tk}{4} \right)^{1/2-\beta'} \left(\frac{tl}{4} \right)^{1/2-\beta''} \times$$

$$I_{\beta-1/2}(th/2) I_{\beta'-1/2}(tk/2) I_{\beta''-1/2}(tl/2) dt, \quad (3.2)$$

iii)

$$a^\alpha \Gamma(-\alpha) F_A^{(3)}(-\alpha, \beta, \beta', \beta'', \gamma, \gamma', \gamma'', h/a, k/a, l/a) \sim$$

$$\Gamma(-\beta-\beta'-\beta''-\alpha) \frac{\Gamma(2\beta)}{\Gamma(\beta)} \frac{\Gamma(2\beta')}{\Gamma(\beta')} \frac{\Gamma(2\beta'')}{\Gamma(\beta'')} h^{-\beta} k^{-\beta'} l^{-\beta''} (a-h-k-l)_+^{\alpha+\beta+\beta'+\beta''} \quad (3.3)$$

as $a \rightarrow 0$.

Proof

i) is a consequence of the formulas

$$a^\alpha F_A^{(3)}(-\alpha, \beta, \beta', \beta'', \gamma, \gamma', \gamma'', h/a, k/a, l/a) =$$

$$\sum_{m,n,p \geq 0} \frac{(-\alpha)_{m+n+p} (\beta)_m (\beta')_n (\beta'')_p}{(\gamma)_m m! (\gamma')_n n! (\gamma'')_p p!} h^m k^n l^p a^{-m-n-p+\alpha} \quad (3.4)$$

$$\frac{d}{da} [a^\alpha F_A^{(3)}](-\alpha, \beta, \beta', \beta'', \gamma, \gamma', \gamma'', h/a, k/a, l/a) = -\alpha a^{\alpha-1} \times$$

$$\sum_{m,n,p \geq 0} \frac{(-\alpha+1)_{m+n+p} (\beta)_m (\beta')_n (\beta'')_p}{(\gamma)_m (\gamma')_n (\gamma'')_p m! n! p!} h^m k^n l^p a^{-m-n-p} \quad (3.5)$$

$$\frac{d}{da} [a^\alpha F_A^{(3)}](-\alpha, \beta, \beta', \gamma, \gamma', \gamma'', h/a, k/a, l/a) = -\alpha a^{\alpha-1} \times$$

$$F_A(-\alpha+1, \beta, \beta', \beta'', \gamma, \gamma', \gamma'', h/a, k/a, l/a) \quad (3.6)$$

To prove ii) we use the formulas ([8], p.237)

$$I_\nu(z) = \frac{2z^\nu e^{-z}}{\sqrt{\pi}\Gamma(\nu+1/2)} \int_0^1 e^{2zt} [t(1-t)]^{\nu-1/2} dt \quad (3.7)$$

where $\alpha \in C, \beta, \gamma, z \in \mathbb{R}^n$ [1], p.115

$$F_A^{(3)}(\alpha, \beta, \gamma, z) = c \int_0^1 \int_0^1 \int_0^1 \prod_{j=1}^{j=3} (1-u_j)^{\gamma_j-\beta_j-1} u_j^{\beta_j-1} \left(1 - \sum_{j=1}^{j=3} u_j z_j\right)^{-\alpha} du_j \quad (3.8)$$

where

$$c = \frac{\prod_{j=1}^{j=3} \Gamma(\gamma_j)}{\prod_{j=1}^{j=3} \Gamma(\beta_j) \Gamma(\gamma_j - \beta_j)} \quad (3.9)$$

iii) The Proof of iii) uses ii) and the formula ([8], p.240)

$$I_{\beta_j-1/2}(x) = \frac{\Gamma(2\beta_j)}{\Gamma(\beta_j)\Gamma(\beta_j+1/2)} 2^{-2\beta_j+1/2} x^{-1/2} (1 + O(|x|^{-1})) \quad (3.10)$$

To finish the proof of the theorem 1.2, we prove the limit conditions in (b): from iii) of the Lemma 2.1 and the Legendre duplication formula for the Γ -Euler function [6], p.3

$$2^{2z-1} \Gamma(z) \Gamma(z+1/2) = \sqrt{\pi} \Gamma(2z). \quad (3.11)$$

we have For $t \rightarrow 0$ and $p, p' \in \mathbb{R}_+^3$

$$W_{(b,b',b'')}^{\mathbb{R}^3}(t, p, p') \sim c_3 1_{\{|p-p'|<t\}}. \quad (3.12)$$

The polar coordinates $p' = p + r\omega$ for $t \rightarrow 0$

$$u(t, p) \sim c_3 \frac{\partial}{\partial t} \int_{\mathbb{R}^3} 1_{\{|p-p'|<t\}} f(p') dp' \quad (3.13)$$

$$u(t, p) \sim c_3 \frac{\partial}{\partial t} \int_0^t f_p^\#(r) r^2 dr \quad (3.14)$$

$$f_p^\#(r) = \frac{1}{6} \int_{S^2} f(p + r\omega) d\omega \quad (3.15)$$

set $r = st$ in the expression above and we see that the limit condition (b) is satisfied and the proof of the theorem B is finished.

References

- [1] P. Appell, Kampe de Fariet, *Fonctions Hypergeometriques et Hyper- spheriques, Polynome d Hermite, Gauthier-Villars, Paris (1926).*
- [2] C.P. Boyer, *Lie theory and separation of variables for the equation $iUt + (\frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2})U = 0$, SIAM J. Math. Anal., 7 (1976), 230-263.*
- [3] Cheeger, J., Taylor, M. *On the diffraction of waves by canonical singularities I, Comm. Pure Appl. Math. 35(3) : 275 – 331, 1982.*
- [4] Exton H., *Multiples hypergeometric functions and Applications, Ellis Horwood Chichester 1976.*
- [5] Folland G. B., *Introduction to partial differential equations, Princeton university press, Princeton N. J. 1976.*
- [6] Magnus F., Oberhettinger and Soni R. P., *Formulas and Theorems for special functions of Mathematical Physics, Third enlarged edition, Springer-Verlag Berlin Heidelberg New York (1966).*
- [7] M.V. Ould Moustapha, *The heat, resolvent and wave kernels with biinverse square potential on the Euclidian plane, Int. Journ. Appl. Math. 27, No 2 (2014), 127-136.*

[8] N.M. Temme, *Special Functions. An Introduction to the Classical Functions of Mathematical Physics*, John Wiley and Sons, Inc., New York- Singapore (1996).

*Department of Mathematics
College of Arts and Sciences at Al Qurayat
Al-Jouf University Kingdom of Saudi Arabia
and
Département de Mathématiques et Informatique
Faculté des Sciences et Techniques
Université de Nouakchott Al-Aasriya
Nouakchott-Mauritanie
Email: mohamedvall.ouldmoustapha230@gmail.com*